

May 13, 2024

New York State Department
of Environmental Conservation
Division of Materials Management
Pesticide Product RegistrationClassified for
"RESTRICTED USE"
in New York State
under 6NYCRR Part 326

DURENTIS™

INSECTICIDE

24(c) Special Local Need Label

**FOR USE IN NASSAU, SUFFOLK,
KINGS OR QUEENS COUNTIES IN
NEW YORK STATE ON TURFGRASS ONLY**

EPA SLN No.:
EPA Reg. No. 279-9674

**FOR DISTRIBUTION AND USE
ONLY IN NEW YORK**

For foliar and systemic control of insect pests in or on golf courses; sports and recreation turf.

This label expires and must not be distributed or used in accordance with this SLN registration after December 31, 2029.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- This state-specific section 24(c) labeling must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, and precautions on the EPA registered label for Durentis Insecticide (EPA Reg. No. 279-9674) and this label.

USE RESTRICTIONS

- **DO NOT** let people and pets enter the treated area until treatment has dried.
- **DO NOT** use for commercial grass seed production.
- **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** apply this product with aerial application equipment.
- **DO NOT** apply more than 6.68 fl oz DURENTIS™ insecticide (equivalent to 0.26 lb of any chlorantraniliprole- containing products) per acre per calendar year to plants produced or grown outdoors.

The following restrictions are required to permit use of this product in the State of New York.

- **Golf course greens and tee boxes:**
 - DO NOT apply this product within 25 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).
- **Golf course fairways:**
 - DO NOT apply this product within 50 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).
- **Other Use Sites:**
 - DO NOT apply this product within 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).

Turfgrass Application rate

Target Pest	Product Application Rate			Application Instructions
	fl. oz product/ 1000 sq ft	lb. ai/ acre	fl. oz product/ acre	
Armyworms Cutworms Sod Webworms	0.016 – 0.031	0.026 – 0.052	0.68-1.33	To ensure optimum control, delay watering (irrigation) or mowing for 24 hours after application. If the grass area is being maintained at a mowing height of greater than 1 inch, then the higher of the listed application rates 0.031 fl oz per 1000 sq. ft. (1.33 fl oz per acre) may be required during periods of high pest pressure.
Billbugs	0.062 – 0.153	0.104 – 0.260	2.68 – 6.68	Apply when overwintered adult billbugs are first observed during April and May. Degree day models have been developed to optimize application timing. Consult your State Cooperative Extension Service for information specific to your region. In temperate regions, spring applications targeting billbug adults will also provide preventative control of white grubs.
White Grubs including: <ul style="list-style-type: none"> • Asian garden beetle • Black Turfgrass Ataenius • Japanese Beetle • European Masked Chafer • Northern Masked Chafer • Southern Masked Chafer • Oriental Beetle • Green June Beetle • May/ June Beetle 	0.062 – 0.123	0.104 – 0.209	2.68 – 5.37	Irrigation of at least 0.1-0.2” is recommended to help move the active ingredient into the soil. Best results will be achieved from preventative applications made just prior to peak activity when grubs are nearest the soil surface. Treatment should be applied in April through early September, with suggested application before June. The higher rate listed may be required [for early-season (April and May) applications where a long residual is required or in late August or early September when less-sensitive mid-instar grubs are present at the time of application. Exact application timing should be confirmed by the local Agricultural Extension Service in your geographic area. Confirmation of the need for treatment may be based on historical records of the site, current adult trap counts or scouting, or other applicable methods.
USE RESTRICTIONS				
<ul style="list-style-type: none"> • Maximum Single Application Rate: <ul style="list-style-type: none"> ○ Do not apply more than 6.68 fl oz of product per acre (0.26 lb ai per acre). • Maximum Annual Application Rate: <ul style="list-style-type: none"> ○ Do not apply more than 6.68 fl oz of product per acre (0.26 lb ai/ acre/year). • Do not enter or allow others to enter treated areas until spray has dried. 				

IMPORTANT NOTICE

BEFORE BUYING OR USING FMC PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY, AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS.

It is a violation of Federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This Bulletin contains supplemental instructions for the use of the reference FMC product. Follow these instructions carefully. This Bulletin must be in the possession of the User at the time of application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using the FMC product(s) covered by this bulletin. FMC will not be responsible for losses or damages resulting from the use of this product in any manner not specifically stated by FMC on the Section 3 label or other labels, or this Bulletin. User assumes all risks associated with any such non-specified use. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability of the Section 3 Federal product label.

FMC, the FMC Logo, and DURENTIS are trademarks of FMC Corporation and/or an affiliate.
© 2023 FMC Corporation. All rights reserved.